

# KPHITE<sup>®</sup> RX

## SYSTEMIC BACTERICIDE

EPA REG. NO. 73806-1

### FIFRA 2(ee) RECOMMENDATION FOR USE IN FLORIDA and TEXAS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency.

**Citrus crops (including subtypes, hybrids and citrus like crops), Avocado and Other Tropical and Subtropical Fruits (Bearing and Non Bearing).**

#### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered labels.

#### GENERAL APPLICATION INSTRUCTIONS

	APPLICATION METHOD	RATE/APPLICATION
Botryosphaeria dothidea Cercospora (Cercospora arachidicola) Cercosporidium (Cercosporidium personatum) Clavibacter (michiganensis subsp. Nebraskensis) Fusarium Phytophthora (soil borne and aerial phases, including brown rot) Pseudomonas spp. (including syringae) Pythium Ralstonia solanacearum Rhizoctonia Sclerotinia Xanthomonas spp. (including campestris) Huanglongbing (HLB, Citrus Greening Disease) <sup>^</sup>	Foliar Spray Application	1-3 quarts in a minimum of 50 gallons of water/acre.
	Aerial Application Equipment	1-3 quarts in a minimum of 5 gallons of water/acre. Not to exceed a 10% concentration.
	Chemigation <sup>^</sup>	4-8 quarts in at least 400 gallons of water/acre.
	Transplant Drench/Dip	2 quarts/100 gallons of water, 2/3 fluid ounce/gallon of water.  Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Trunk or Bark Injection	3 teaspoons/linear yard of canopy width or 2 inches of trunk diameter at breast height (DBH). Inject directly into the tree.  Apply according to injection equipment instructions. Adjust rate on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
	Soil /Root Drench <sup>^</sup>	1-2 gallons/100 gallons of water or 1.25-2.5 fluid ounces per gallon.  Apply 5-10 gallons of diluted solution per tree. Adjust volume as needed to thoroughly drench area around trunk. Apply at 3-4 week intervals and repeat as needed.
Alternaria Anthracnose Downy Mildew Powdery Mildew Rust Huanglongbing (HLB, Citrus Greening Disease) <sup>***</sup>	Soil Directed Application <sup>^</sup>	4-8 quarts in min 20 gallons of water/acre.
	Foliar Spray Application	1-4 quarts in a minimum of 100 gallons of water/acre.
	Aerial Application Equipment	1-4 quarts in a minimum of 5 gallons of water/acre. Not to exceed a 10% concentration.

\*\*\* Use a minimum of 2 quarts per acre. The addition of a Fertilizer, such as Renew<sup>®</sup>RX or Keypon RX, may improve control.

A non-ionic surfactant may also be used. <sup>^</sup> For use on indicated application methods only.

**All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.** This FIFRA Section 2(ee) recommendation contains new or additional Diseases, which may not appear on the package label. Read and observe the cautionary statements plus all other information appearing on the product label.

#### REGISTRANT : PLANT FOOD SYSTEMS, INC

P.O. BOX 775, 2827 Union Street, Zellwood, FL. 32798  
1 (800) 343-7775  
KPHITE.com

